

## **SPRINGBORO PEDIATRICS, INC.**

8 Sycamore Creek Drive  
Springboro, Ohio 45066  
Phone: 937 748-KIDS (5437)  
Fax: 937 748-5434

Brian Esselstein, M.D.

Charles Hutchison, M.D.

Nalini Aggarwal, M.D.

### **IMMUNIZATION/PHYSICAL EXAM SCHEDULE**

2 MONTHS	Well child check *Pentacel (combination vaccine - DTaP/Hib/IPV) Hep B Pneumovax (Pneumococcal vaccine) Rotavirus (initial vaccine has to be given at 2 months)
4 MONTHS	Well child check *Pentacel (combination vaccine - DTaP/Hib/IPV) Hep B (given only if not received at birth or at 2 weeks) Pneumovax (Pneumococcal vaccine) Rotavirus (only if given at 2 months)
6 MONTHS	Well child check *Pentacel (combination vaccine - DTaP/Hib/IPV) Hep B Pneumovax (Pneumococcal vaccine) Rotavirus (only if given at 2 & 4 months)
9 MONTHS	Well child check only – no shots
12 MONTHS	Well child check MMR #1 (Measles, Mumps, Rubella vaccine) **Varivax (Chicken pox vaccine) Pneumovax (Pneumococcal vaccine)
15 MONTHS	Well child check *Pentacel (combination vaccine - DTaP/Hib/IPV)
18 MONTHS	Well child check

2 YEARS - 4 YEARS	Yearly well child checks – no shots
5 YEARS	Well child check DTaP (Diphtheria, Tetanus, Acellular Pertussis) IPV (Poliovirus vaccine) MMR #2 (Measles, Mumps, Rubella vaccine) If not received previously **Varivax Booster is recommended
6 YEARS - 18 YEARS	Physical exam every 2 years, until in Junior High/High School – yearly exam at that time if in sports and/or marching band which requires a yearly physical
12 YEARS - 15 YEAR	****Tdap Adult (Tetanus, Diphtheria, Acellular Pertussis) <b>AND</b> Menactra (Meningitis)
9 YEARS - 26 YEARS	HPV (Human Papillomavirus Vaccine) recommended for girls (3 doses) available for boys (3 doses)

Flu vaccine injection is recommended yearly for children 6 months and older  
**OR**  
Flumist nasal spray vaccine is available yearly for children 2 years and older

\*Pentacel (DTaP - Diphtheria, Tetanus, Acellular Pertussis), (Hib – Haemophilus Influenzae),  
(IPV – Polio)

\*\*Varivax is a chicken pox vaccine that can be given any time after 12 months of age

\*\*\* DTaP (Diphtheria, Tetanus, Acellular Pertussis), Hib (Haemophilus Influenzae Type b)

\*\*\*\*Tdap should be given every 7-10 years